

Nonholonomic constraints and Voronec's equations

Nivaldo A. Lemos

Departamento de Física

Universidade Federal Fluminense

Av. Litorânea s/n, Boa Viagem

24210-340 Niterói, RJ

Brazil

February 2, 2008

Abstract

Is it allowed, in the context of the Lagrange multiplier formalism, to assume that nonholonomic constraints are already in effect while setting up Lagrange's function? This procedure is successfully applied in a recent book [L. N. Hand and J. D. Finch, *Analytical Mechanics*] to the problem of the rolling penny, but it does not work in general, as we show by means of a counterexample. It turns out that in many cases the use of nonholonomic constraints in the process of construction of the Lagrangian is allowed, but then the correct equations of motion are the little known Voronec's equations.

In the Lagrange multiplier formalism as applied to nonholonomic systems, the Lagrangian is written as if there were no constraints. The nonholonomic constraints are taken into account in the formulation of the equations of motion, but not during the construction of the Lagrangian. Setting up the Lagrangian assuming that the constraints are already in effect is completely equivalent to substituting the constraint equations into the Lagrangian written as if there were no constraints. It is tempting to take it for granted that the ensuing reduced Lagrangian together with the relevant constraint equations always lead to the correct equations of motion for the system. The procedure just described is successfully used in a recent book [1] to solve the problem of a penny rolling on an inclined plane. Unfortunately, contrary to what the mentioned book appears to suggest, this approach is not valid in general, as we proceed to show with the help of a counterexample.

Consider a homogeneous sphere rolling without slipping on a horizontal plane. This problem is treated by the Lagrange multiplier method in [2]. Let X, Y, Z be cartesian axes fixed in space with the Z -axis perpendicular to the plane. The principal moments of inertia with respect to the center of the sphere are all equal to $2mR^2/5$. With x, y the coordinates of the center of the sphere, the Lagrangian, being equal to the kinetic energy, is given by

$$L = \frac{m}{2}(\dot{x}^2 + \dot{y}^2) + \frac{mR^2}{5}\omega^2 . \quad (1)$$

The constraint equations are

$$\dot{x} = R\omega_y = R\dot{\theta} \sin \phi - R\dot{\psi} \sin \theta \cos \phi , \quad \dot{y} = -R\omega_x = -R\dot{\theta} \cos \phi - R\dot{\psi} \sin \theta \sin \phi . \quad (2)$$

In terms of the Euler angles ϕ, θ, ψ the Lagrangian (1) takes the form

$$L = \frac{m}{2}(\dot{x}^2 + \dot{y}^2) + \frac{mR^2}{5}(\dot{\phi}^2 + \dot{\theta}^2 + \dot{\psi}^2 + 2\dot{\phi}\dot{\psi} \cos \theta) . \quad (3)$$

According to the method employed in [1], which takes into account the rolling constraint in the construction of the Lagrangian, the kinetic energy is written in terms of the rotational degrees of freedom alone by taking the moments of inertia with respect to the contact point of the sphere with the plane. The Lagrangian is now

$$\bar{L} = \frac{1}{2} \frac{7mR^2}{5} (\omega_x^2 + \omega_y^2) + \frac{1}{2} \frac{2mR^2}{5} \omega_z^2 . \quad (4)$$

This is exactly the Lagrangian that one obtains by inserting the constraint equations (2) into the Lagrangian (1), which was written as if there were no constraints. In terms of the Euler angles the

reduced Lagrangian (4) becomes

$$\bar{L} = \frac{7mR^2}{10}(\dot{\theta}^2 + \dot{\psi}^2 \sin^2 \theta) + \frac{mR^2}{5}(\dot{\phi}^2 + \dot{\psi}^2 \cos^2 \theta + 2\dot{\phi}\dot{\psi} \cos \theta) . \quad (5)$$

Since the variables x and y do not appear in $\bar{L}$, according to the reasoning in [1] the constraint equations (2) are no longer relevant to the formulation of the equations of motion. In particular, the Lagrange equation for θ is

$$\frac{d}{dt} \left(\frac{\partial \bar{L}}{\partial \dot{\theta}} \right) - \frac{\partial \bar{L}}{\partial \theta} = 0 \implies 7\ddot{\theta} - 5\dot{\psi}^2 \sin \theta \cos \theta + 2\dot{\phi}\dot{\psi} \sin \theta = 0 . \quad (6)$$

The treatment of this problem by the Lagrange multiplier method shows that the two Lagrange multipliers vanish [2]. According to equation (58d) of [2] the correct equation of motion for θ is

$$\ddot{\theta} + \dot{\phi}\dot{\psi} \sin \theta = 0 , \quad (7)$$

which is completely different from equation (6). The differential equations (6) and (7) generally yield different solutions for θ because $\phi, \theta, \psi, \dot{\phi}, \dot{\theta}, \dot{\psi}$ can be arbitrarily chosen at any particular instant t_0 . It is clear, therefore, that the approach suggested in [1] lacks generality, since it works for the rolling penny but fails for the rolling sphere.

In a previous paper [3] we remarked that it is possible to perform a reduction of the Lagrangian taking into account the constraints, but in this case the correct equations of motion are Voronec's equations. Given a dynamical system described by the configuration variables $q_1, \dots, q_n$, suppose the first m velocities are independent and the $k = n - m$ remaining velocities can be expressed in terms of the independent ones by means of the equations

$$\dot{q}_{m+l} - \sum_{j=1}^m a_{lj} \dot{q}_j = 0 , \quad l = 1, \dots, k , \quad (8)$$

where the coefficients a_{lj} are functions of the generalized coordinates $q_1, \dots, q_n$.

Let L be the Lagrangian written without taking into account the nonholonomic constraint equations (8). If the last k velocities are eliminated from the Lagrangian by means of equations (8), a reduced Lagrangian $\bar{L}$ results:

$$L(q_1, \dots, q_n, \dot{q}_1, \dots, \dot{q}_n, t) = \bar{L}(q_1, \dots, q_n, \dot{q}_1, \dots, \dot{q}_m, t) . \quad (9)$$

Voronec's equations of motion are [3, 4]

$$\frac{d}{dt} \left(\frac{\partial \bar{L}}{\partial \dot{q}_i} \right) - \frac{\partial \bar{L}}{\partial q_i} = \sum_{\nu=1}^k \frac{\partial \bar{L}}{\partial q_{m+\nu}} a_{\nu i} + \sum_{\nu=1}^k \sum_{j=1}^m \frac{\partial L}{\partial \dot{q}_{m+\nu}} b_{ij}^{\nu} \dot{q}_j , \quad i = 1, \dots, m , \quad (10)$$

where

$$b_{ij}^{\nu} = \frac{\partial a_{\nu i}}{\partial q_j} - \frac{\partial a_{\nu j}}{\partial q_i} + \sum_{\mu=1}^k \left(\frac{\partial a_{\nu i}}{\partial q_{m+\mu}} a_{\mu j} - \frac{\partial a_{\nu j}}{\partial q_{m+\mu}} a_{\mu i} \right) . \quad (11)$$

Setting $q_1 = \phi, q_2 = \theta, q_3 = \psi, q_4 = x, q_5 = y$, in the present case $m = 3$ and $k = 2$. The constraint equations (2) can be written in the form (8) with

$$a_{11} = 0, a_{12} = R \sin \phi, a_{13} = -R \sin \theta \cos \phi, a_{21} = 0, a_{22} = -R \cos \phi, a_{23} = -R \sin \theta \sin \phi . \quad (12)$$

The definition (11) furnishes immediately the only nonvanishing coefficients b_{ij}^{ν} :

$$b_{12}^1 = -b_{21}^1 = -R \cos \phi, \quad b_{13}^1 = -b_{31}^1 = -R \sin \theta \sin \phi, \quad b_{23}^1 = -b_{32}^1 = R \cos \theta \cos \phi, \quad (13)$$

$$b_{12}^2 = -b_{21}^2 = -R \sin \phi, \quad b_{13}^2 = -b_{31}^2 = R \sin \theta \cos \phi, \quad b_{23}^2 = -b_{32}^2 = R \cos \theta \sin \phi . \quad (14)$$

It follows that Voronec's equation for θ is

$$\frac{7mR^2}{5} \ddot{\theta} - mR^2 \dot{\psi}^2 \sin \theta \cos \theta + \frac{2mR^2}{5} \dot{\phi} \dot{\psi} \sin \theta = m\dot{x}(b_{21}^1 \dot{\phi} + b_{23}^1 \dot{\psi}) + m\dot{y}(b_{21}^2 \dot{\phi} + b_{23}^2 \dot{\psi}) . \quad (15)$$

The use of the constraint equations (2) and a little algebra reduce the above equation to

$$\ddot{\theta} + \dot{\phi} \dot{\psi} \sin \theta = 0 , \quad (16)$$

which coincides with the correct equation for θ furnished by the Lagrange multiplier method. It is equally straightforward to check that the remaining Voronec equations for ϕ and ψ coincide with those obtained by the Lagrange multiplier method.

In short, the method employed in [1] is not valid in general and should not be taught to students. The correct result obtained for the rolling penny is the product of a mere accident.

As a general rule, the use of nonholonomic constraints while setting up the Lagrangian is allowed, but the correct equations of motion are Voronec's equations, and not the ones given by the Lagrange multiplier method.

References

- [1] L. N. Hand and J. D. Finch, *Analytical Mechanics* (Cambridge University Press, Cambridge, 1998), Sec. 2.8.
- [2] E. J. Saletan and A. H. Cromer, *Theoretical Mechanics* (John Wiley and Sons, New York, 1971), Sec. V-4 (e).
- [3] N. A. Lemos, Rev. Bras. Ens. Fís. **25**, 28 (2003).
- [4] Ju. I. Neĭmark and N. A. Fufaev, *Dynamics of Nonholonomic Systems* (American Mathematical Society, Providence, Rhode Island, 1972), Chap. III, Sec. 3.